# Social Learning

Duarte Gonçalves
University College London

Topics in Economic Theory

## Overview

- 1. Social Learning: Checking Neighbours' Crop Yields
- 2. Warm-Up: Gaussian-Quadratic Model
- 3. Information Cascades
- 4. Further Topics

## Overview

- 1. Social Learning: Checking Neighbours' Crop Yields
- 2. Warm-Up: Gaussian-Quadratic Model
- Information Cascades
- Further Topics

## Social Learning

Numerous situations in which we try to learn from others' actions

Knowing a fim is successful is informative about its quality and genre (unlikely to be awful, but also unlikely to be an indie film)

Seeing price of a stock going up: infer more positive outlook on firm's fundamentals more likely.

Product purchases, migration decisions, bachelor choices, technology adoption, take-up of microfinance, etc.

Can the crowd get it wrong? When?

## Setup

## **Basic Ingredients**

 $\theta \in \Theta$  unknown parameter;  $\theta \sim P_{\Theta}$ .

Action: Each agent *n* takes action  $a_n \in A \subseteq \mathbb{R}$ .

Public History:  $h_t := \{a_\ell, \ell < t\}, h_0 = \emptyset$ .

Payoffs:  $u(a_n, \theta, r_n)$ ; Type: Agent *n*'s type  $r_n$ ; *R* types.

Private Information: Each agent n has private information about  $\theta$ .

Important how large A; coarser set restricts how much info conveyed.

## **Examples**

**Binary Actions:**  $\Theta = \{H, L\}$ ,  $A = \{0, 1\}$ , e.g. buy/invest/adopt/migrate/etc vs don't. WLOG  $u(a, \theta, r) := a(1_{\{\theta = H\}} - r)$ .

**Gaussian-Quadratic:**  $\Theta = A = \mathbb{R}$ , e.g., predict variable:  $u(a, \theta, r) := -(\theta - a)^2$  or investment problem  $u(a, \theta, r) := 2\theta a - a^2$ .

#### **Private and Public Information:**

```
Private Signal: s_n \sim P_S(\cdot \mid \theta) iid (conditional on \theta); p_n^0(\theta) \propto p_0(\theta) P_S(s_n \mid \theta). p_n^0 = f_p(s_n), s_n \sim P_S(\cdot \mid \theta); hence p_n^0 \sim P_P(\cdot \mid \theta) for some P_P s.t. \mathbb{E}[p_n^0] = P_{\theta}. Private Belief: p_n^0 \sim P_P.

Public Belief: q_t(\theta) = \mathbb{P}(\theta \mid h_t). \pi(h_t \mid \theta) := \mathbb{P}(h_t \mid \theta); q_t(\theta) \propto p_0(\theta)\pi(h_t \mid \theta).

Private Posterior Belief: Private belief + updating based on public history. p_t(\theta) \propto p_0(\theta) P_S(s_t \mid \theta)\pi(h_t \mid \theta) = q_t(\theta) p_t^0(\theta)/p_0(\theta). Updating mapping BU: (p_t^0, q_t) \mapsto BU(p_t^0, q_t) = p_t.
```

## Timing: Within each period t:

- 1. Agent *t* observes private signal  $s_n$  and forms interim private belief  $p_t^0$ .
- 2. Agent observes public history  $h_t = (a_\ell, \ell < t)$  and forms private posterior belief  $p_t$ .
- 3. Agent t takes action  $a_t$ .

wPBE (focusing on on-path)

## Herd, Cascade

## **Definition**

- (i) Actions converge if  $\sum_t a_t/t \to a_\infty$  for some  $a_\infty \in A$ .
- (ii) A herd occurs at t if, given the public belief, agent t's optimal action is the same as agent (t-1)'s.
- (iii) A herd starts at T if it occurs at any  $t \geq T$ .
- (iv) A cascade occurs at *t* if, given the public belief, agent *t*'s optimal action is a.s. independent of *t* private signal.
- (v) A cascade starts at T if it occurs at any  $t \geq T$ .
- (vi) A cascade that occurs at  $\infty$  is a limit cascade.

#### Remark

Cascade in finite time  $\implies$  Limit cascade, Herd in finite time  $\implies$  Actions converge. If A finite, Limit cascade  $\implies$  Herd in finite time.

## Overview

- Social Learning: Checking Neighbours' Crop Yields
- 2. Warm-Up: Gaussian-Quadratic Model
- 3. Information Cascades
- Further Topics

## Warm-Up: Gaussian-Quadratic Model

$$u(a,\theta) = -(\theta - a)^2$$
;  $\theta \sim N(\theta_0, \rho_{\theta}^{-1})$ ;  $s_t = \theta + \varepsilon_t$ ;  $\varepsilon_t \sim N(0, \rho_{\varepsilon}^{-1})$  iid; indep. from  $\theta$ . Public Belief:  $N(\mu_t, \rho_t^{-1})$ .

Posterior Belief = Private Signal + Public Belief: 
$$N(\tilde{\mu}_t, \tilde{\rho}_t^{-1})$$

$$\tilde{\mu}_t = (1 - w_t)\mu_t + w_t s_t; \quad \tilde{\rho}_t = \rho_t + \rho_\epsilon; \quad w_t = \rho_\epsilon/\tilde{\rho}_t.$$

Conditional on posterior belief,  $a_t = \arg\max_a \mathbb{E}[u(a, \theta) \mid h_t, s_t] = \mathbb{E}[\theta \mid h_t, s_t] = \tilde{\mu}_t$ .

Social Learning: As 
$$\mu_t$$
,  $a_t$ ,  $w_t$  are known, agent  $t+1$  can infer  $s_t = (a_t - (1-w_t)\mu_t)/w_t$   
 $\implies \mu_{t+1} = \tilde{\mu}_t$  and  $\rho_{t+1} = \tilde{\rho}_t$ .

As 
$$a_t = \mu_t = \frac{\rho_\theta}{\rho_t} \theta_0 + \frac{t\rho_\epsilon}{\rho_t} \sum_{\ell < t} s_\ell \to \theta$$
 a.s.  $\Longrightarrow$  learning is complete.

No herd, no cascade in finite time.

Imitation:  $\mathbb{E}[|a_t - a_{t-1}|] = w_t \mathbb{E}[|(s_t - a_{t-1})|] \to \mathbf{0}$ : actions more similar over time.

Heterogeneity:  $u(a, \theta, r) = -(\theta + r - a)^2$ ,  $r \sim N(0, \rho_r^{-1})$ ; results hold, but learning slower.

Gonçalves (UCL) Social Learning

# Warm-Up: Gaussian-Quadratic Model

Costly signal: 
$$u(a, \theta) = -(\theta - a)^2 - c(\rho_{\varepsilon})$$
;  $c \ge 0$ ,  $c' > 0$ ,  $c'' > 0$ .

Claim: Social learning stops in finite time, i.e.,

$$\exists T < \infty : \forall t > T, \ \rho_t = \rho_T \iff \rho_{\varepsilon,t} = 0$$

- $\mathbb{E}[u(a_t, \theta) \mid h_t, s_t] = -\mathbb{E}[(\theta \mathbb{E}[\theta \mid h_t, s_t])^2 \mid h_t, s_t] c(\rho_{\epsilon,t}) = -(\rho_t + \rho_{\epsilon,t})^{-1} c(\rho_{\epsilon,t}).$ FOC:  $(\rho_t + \rho_{\epsilon,t})^{-2} - c'(\rho_{\epsilon,t}) = 0.$
- When  $\rho_t \geq c'(0)^{1/2} \implies c'(\rho_{\epsilon,t}) \geq c'(0) \geq \rho_t^{-2} \geq (\rho_t + \rho_{\epsilon,t})^{-2} \implies \rho_{\epsilon,t} = 0.$
- $c'(\rho_{\varepsilon,t-1}) = (\rho_{t-1} + \rho_{\varepsilon,t-1})^{-2} = \rho_t^{-2} \ge (\rho_t + \rho_{\varepsilon,t})^{-2} = c'(\rho_{\varepsilon,t})$  $\Longrightarrow \rho_{\varepsilon,t}$  decreasing in t.

# Warm-Up: Gaussian-Quadratic Model

Costly signal: 
$$u(a, \theta) = -(\theta - a)^2 - c(\rho_{\varepsilon})$$
;  $c \ge 0$ ,  $c' > 0$ ,  $c'' > 0$ .

Claim: Social learning stops in finite time, i.e.,

$$\exists T < \infty : \forall t > T, \ \rho_t = \rho_T \iff \rho_{\varepsilon,t} = 0$$

$$FOC: (\rho_t + \rho_{\varepsilon,t})^{-2} - c'(\rho_{\varepsilon,t}) = 0.$$

 $\implies \rho_{\epsilon,t}$  decreasing in t.

• Assume  $\rho_{\epsilon,t} > 0$ .

$$\Rightarrow c'(0)^{-1/2} \geq c'(\rho_{\epsilon,t})^{-1/2} \qquad \qquad \rho_{\epsilon,t} \text{ decreasing in } t; c'' > 0$$

$$= \rho_t + \rho_{\epsilon,t} \qquad \qquad \text{FOC, } 0 < \rho_{\epsilon,t}$$

$$= \rho_{t-1} + \rho_{\epsilon,t-1} + \rho_{\epsilon,t} = \rho_0 + \rho_{\epsilon,1} + \dots + \rho_{\epsilon,t}$$

$$= \rho_0 + \sum_{\ell=0}^{t-1} c'(\rho_{\epsilon,t-\ell})^{-1/2} \qquad \qquad \text{FOC, } 0 < \rho_{\epsilon,t} \leq \rho_{\epsilon,\ell} \ \forall \ell \leq t$$

$$\geq tc'(\rho_{\epsilon,1})^{-1/2} \qquad \qquad \rho_{\epsilon,t} \text{ decreasing in } t; c'' > 0$$

$$\Rightarrow t < (c'(\rho_{\epsilon,1})/c'(0))^{1/2}.$$

Gonçalves (UCL) Social Learning

## Crashes and Booms

Adjustment to economic shock tends to be first slow and then abrupt (e.g., a market crash). Why?

**Model sketch:** population every period takes irreversible investment decisions; start with high public confidence in 'good times' but state is bad.

Sigmoid-shaped adjustment of beliefs from social learning to change in state features:

- (1) slow evolution (somewhat business as usual),
- (2) fast adjustment to new reality (negative shock), and
- (3) stabilising around 'new normal'.

Caplin and Leahy (1994 AER); Chamley (2010 Ch. 4.6).

Fast changing state would temper these abrupt changes.

## Overview

- 1. Social Learning: Checking Neighbours' Crop Yields
- 2. Warm-Up: Gaussian-Quadratic Mode
- 3. Information Cascades
  - Setup
  - Social Learning
  - Cascades
  - Robustness of Cascades
- 4. Further Topics

## Setup

Version of Bikchandani, Hirschleifer, and Welch (1992 JPE). (but using martingale tools employed by Smith and Sørensen (2000 Ecta).)

### Setup

together.

State:  $\theta \in \Theta = \{H, L\}, p_0 = \mathbb{P}(\theta = H) \in (0, 1).$ 

Signal:  $s_n \in S = \{1, 2, ..., N_S\}$ ;  $s_n \sim P_S(\cdot \mid \theta)$ . Assume  $P_S(s \mid \theta) > 0 \quad \forall s, \theta$ . (No fully revealing signals with positive probability.)

Order signals:  $\frac{P_S(1|H)}{P_S(1|L)} < \cdots < \frac{P_S(N_S|H)}{P_S(N_S|L)}$ .

MLRP WLOG when  $|\Theta| = 2$ ; strict MLRP: bundle signals with same likelihood

Payoffs:  $u(a, \theta, r) := a(\mathbf{1}_{\{\theta = H\}} - r)$ . Tie-breaking: Invest iff belief  $\theta = H > r$ .

## Setup

### **Cascade Sets and Other Definitions**

Public History: 
$$h_t = (a_\ell)_{\ell < t}$$
;  $h_0 = \emptyset$ .

Public Belief: 
$$q_t = \frac{p_0 \pi(h_t | H)}{p_0 \pi(h_t | H) + (1 - p_0) \pi(h_t | L)}$$
.

Private Interim Belief: 
$$p_t^0(s) = \frac{p_0 P_S(s|H)}{p_0 P_S(s|H) + (1-p_0) P_S(s|L)}$$
.  $p_t^0 \sim P_P$ .

Let  $\underline{b}$  := inf supp( $P_P$ ) > 0 and  $\bar{b}$  := sup supp( $P_P$ ) < 1. (Recall: finite S.)

### **Private Posterior Belief:**

$$p_t(s) = \frac{q_t P_S(s|H)}{q_t P_S(s|H) + (1-q_t) P_S(s|L)} = \frac{q_t p_t^0/p_0}{q_t p_t^0/p_0 + (1-q_t) (1-p_t^0)/(1-p_0)} =: BU(p_t^0, q_t).$$

Action a's Basin: 
$$I_a' := \{ p \in [0,1] \mid a = \min(\arg\max_{a'} p \, u(a', H, r) + (1-p) \, u(a', L, r)) \}.$$

$$\exists 0 = \hat{\rho}_0 < \hat{\rho}_1 < \hat{\rho}_2 = 1 : l_0^r = [\hat{\rho}_0, \hat{\rho}_1], \, l_1^r = (\hat{\rho}_1, \hat{\rho}_2].$$

(Argument extends beyond binary actions)

Cascade Set for a: set of public beliefs at which agent chooses a regardless of private information.

$$J_a^r := \{q \mid BU(p,q) \in I_a^r, \forall p \in \text{supp}(P_P)\}.$$

$$\exists 0 = \hat{q}_0^{r,-} < \hat{q}_0^{r,+} < \hat{q}_1^{r,-} < \hat{q}_1^{r,+} = 1: J_0^r = [\hat{q}_0^{r,-}, \hat{q}_0^{r,+}] \text{ and } J_1^r = (\hat{q}_1^{r,-}, \hat{q}_1^{r,+}],$$
 with  $BU(b, \hat{q}_2^{r,-}) = \hat{p}_2^r$  and  $BU(\bar{b}, \hat{q}_2^{r,+}) = \hat{p}_{2+1}^r$ .

$$q \in \mathcal{J}_a^r \iff \hat{p}_a \leq BU(\underline{b},q) < BU(p_0,q) = q < BU(\bar{b},q) \leq \hat{p}_{a+1} \implies \mathcal{J}_a^r \subseteq \operatorname{int}(I_a^r).$$

Define  $J^r := \bigcup_a J_a^r$ .

## Not Socially Learning the Wrong Thing

### **Proposition**

In any equilibrium, a fully wrong information cascade a.s. never starts.

### **Proof**

If  $q_T \in \mathcal{J}^r$ , then at any  $t \geq T$ , private signals are ignored in favour of public history. Need to show wp1  $\exists T < \infty : \forall t > Tq_T \in \mathcal{J}^r = [0, \hat{q}_0^+] \cup (\hat{q}_1^-, 1]$ .

Public Likelihood Ratio:  $I(q_t) = (1 - q_t)/q_t$ .

Claim: Conditional on  $\theta = H$ ,  $I(q_t)$  is a martingale.

$$\begin{split} \mathbb{E}[I(q_{t+1}) \mid h_t, \theta = H] &= \mathbb{E}\left[\frac{(1 - p_0)\pi^L(h_{t+1})}{p_0\pi^H(h_{t+1})} \middle| h_t, \theta = H\right] \\ &= \mathbb{E}\left[\frac{\mathbb{P}(a_t \mid h_t, \theta = L)(1 - p_0)\pi^L(h_t)}{\mathbb{P}(a_t \mid h_t, \theta = H)p_0\pi^H(h_t)} \middle| h_t, \theta = H\right] \\ &= I(q_t) \sum_{a} \mathbb{P}(a_t = a \mid h_t, \theta = H) \frac{\mathbb{P}(a_t = a \mid h_t, \theta = L)}{\mathbb{P}(a_t = a \mid h_t, \theta = H)} = I(q_t) \end{split}$$

Note:  $q_t$  is a martingale; conditional on  $\theta = H$  it is a submartingale.

## Not Socially Learning the Wrong Thing

### **Proposition**

In any equilibrium, a fully wrong information cascade a.s. never starts.

## **Proof**

If  $q_T \in J^r$ , then at any  $t \ge T$ , private signals are ignored in favour of public history. Need to show wp1  $\exists T < \infty : \forall t > Tq_T \in J^r = [0, \hat{q}_1^+] \cup (\hat{q}_1^-, 1]$ .

Public Likelihood Ratio:  $I(q_t) = (1 - q_t)/q_t$ .


Claim: Conditional on  $\theta = H$ ,  $I(q_t)$  is a martingale.

$$\begin{split} \mathbb{E}[I(q_{t+1}) \mid h_t, \theta = H] &= \mathbb{E}\left[\frac{(1-p_0)\pi^L(h_{t+1})}{p_0\pi^H(h_{t+1})} \middle| h_t, \theta = H\right] \\ &= \mathbb{E}\left[\frac{\mathbb{P}(a_t \mid h_t, \theta = L)(1-p_0)\pi^L(h_t)}{\mathbb{P}(a_t \mid h_t, \theta = H)p_0\pi^H(h_t)} \middle| h_t, \theta = H\right] \\ &= I(q_t) \sum_{\alpha} \mathbb{P}(a_t = a \mid h_t, \theta = H) \frac{\mathbb{P}(a_t = a \mid h_t, \theta = L)}{\mathbb{P}(a_t = a \mid h_t, \theta = H)} = I(q_t) \end{split}$$

 $\mathit{I}(q_t) \geq$  0. By Doob's martingale convergence theorem, conditional on  $\theta$  =  $\mathit{H}$ ,  $\mathit{I}(q_t) \rightarrow$ 

#### **Theorem**

In any equilibrium, an information cascade starts in finite time a.s.



**Figure 4.2 Representation of a cascade.** In each period, the middle of the vertical segment is the public belief; the top and the bottom of the segment are the beliefs of an optimist (with a private signal s=1) and of a pessimist (with signal s=0). The private signals are  $s_1=0$ ,  $s_2=1$ ,  $s_3=0$ ,  $s_4=1$ ,  $s_5=1$ .

#### Theorem

In any equilibrium, an information cascade starts in finite time a.s.

## **Proof**

Claim:  $supp(q_{\infty}) \subseteq [0, \hat{q}_{0}^{+}] \cup [\hat{q}_{1}^{-}, 1].$ 

Suppose  $q^* \in (\hat{q}_0^+, \hat{q}_1^-) \cap \text{supp}(q_t) \implies \forall \epsilon > 0$ ,  $\mathbb{P}(q_\infty \in (q^* - \epsilon, q^* + \epsilon)) > 0$ .

Hence wp>0  $\exists T < \infty : \forall t \geq T, q_t \in (q^* - \varepsilon, q^* + \varepsilon).$ 

Take  $\varepsilon$  arbitrarily small.  $(q^* - \varepsilon, q^* + \varepsilon) \subset (\hat{q}_0^+, \hat{q}_1^-)$ , an active learning region.

In the active learning region, prob agent takes 0 or 1 is different depending on the true state, and so  $\mathbb{P}(a_t = 1 \mid q_t, \theta = H) \neq \mathbb{P}(a_t = 1 \mid q_t, \theta = H)$ .

$$a_t = 1 \implies q_{t+1} > q_t \text{ and } a_t = 0 \implies q_{t+1} < q_t.$$

Changes are discrete, so can find small  $\varepsilon > 0$  s.t. exit  $(\hat{q}_0^+, \hat{q}_1^-)$  wp>0 starting from  $q_t \in (\hat{q}_0^+, \hat{q}_1^-)$ . Contradiction.

#### Theorem

In any equilibrium, an information cascade starts in finite time a.s.

## **Proof**

Claim: 
$$\mathbb{P}(q_{\infty} = \hat{q}_1^-) = \mathbf{0}$$
.

Suppose  $\mathbb{P}(q_{\infty} = \hat{q}_{1}^{-}) > 0$ .

Hence for positive prob sample of paths  $q_t \uparrow \hat{q}_1^-$ .

 $\implies$  for any  $\varepsilon > 0$ , wp>0  $\exists T < \infty : \forall t \geq T, q_t \in (\hat{q}_1^- - \varepsilon, \hat{q}_1^-].$ 

Note  $a_t = f(s_t, q_t)$ , hence  $supp(q_{t+1} \mid q_t = q)$  is independent from t.

 $\implies q_{t+1}$  is time-homogeneous Markov process.

For small enough  $\varepsilon$ , one has  $\forall q_t \in (\hat{q}_1^- - \varepsilon, \hat{q}_1^-] \implies (a_t = 0 \iff s_t = 1)$ .

#### **Theorem**

In any equilibrium, an information cascade starts in finite time a.s.

## **Proof**

Claim: 
$$\mathbb{P}(q_{\infty}=\hat{q}_{1}^{-})=\mathbf{0}$$
. For small enough  $\boldsymbol{\varepsilon}$ , one has  $\forall q_{t}\in(\hat{q}_{1}^{-}-\boldsymbol{\varepsilon},\hat{q}_{1}^{-}]\Longrightarrow(a_{t}=\mathbf{0}\iff s_{t}=\mathbf{1})$ . Note that, since  $q_{t}$  is a martingale,  $\mathbb{E}[q_{t+1}\mid q_{t}]=q_{t}$  and then  $\frac{q_{t}P_{S}(1|H)}{q_{t}P_{S}(1|H)+(1-q_{t})P_{S}(1|L)}< q_{t}<\frac{q_{t}(1-P_{S}(1|H))}{q_{t}(1-P_{S}(1|H))+(1-q_{t})(1-P_{S}(1|L))}$ . Choose  $\boldsymbol{\varepsilon}>\mathbf{0}$  such that 
$$\frac{\hat{q}_{1}P_{S}(1|H)}{\hat{q}_{1}P_{S}(1|H)+(1-\hat{q}_{1}^{-})P_{S}(1|L)}<\hat{q}_{1}^{-}-\boldsymbol{\varepsilon}<\hat{q}_{1}^{-}<\frac{(\hat{q}_{1}^{-}-\boldsymbol{\varepsilon})(1-P_{S}(1|H))}{(\hat{q}_{1}^{-}-\boldsymbol{\varepsilon})(1-P_{S}(1|H))+(1-(\hat{q}_{1}^{-}-\boldsymbol{\varepsilon}))(1-P_{S}(1|L))}.$$
 Then,  $q_{t}\in(\hat{q}_{1}^{-}-\boldsymbol{\varepsilon},\hat{q}_{1}^{-}]$   $\Longrightarrow q_{t+1}\notin(\hat{q}_{1}^{-}-\boldsymbol{\varepsilon},\hat{q}_{1}^{-}]$ . Contradiction.

#### Theorem

In any equilibrium, an information cascade starts in finite time a.s.

## **Proof**

We have  $\operatorname{supp}(q_{\infty}) \subseteq J^r$ . It remains to be shown that a.s.  $\exists T < \infty : \forall t \geq T, q_t \in J^r$ .

Note  $J^r = \operatorname{int} J^r \cup \{\hat{q}_0^+\}.$ 

 $\text{On } \{q_{\infty} \in \text{int } \mathcal{J}'\}, \\ \text{lim}_{t \to \infty} \ q_t = q_{\infty} \in \text{int } \mathcal{J}' \implies \exists \mathcal{T} < \infty : \forall t \geq \mathcal{T}, \\ q_t \in \mathcal{J}'.$ 

WTS  $\{q_{\infty} = \hat{q}_{\mathbf{0}}^{+}\}$ , a.s.  $\exists T < \infty : \forall t \geq T$ ,  $q_{t} = \hat{q}_{\mathbf{0}}^{+}$ .

#### **Theorem**

In any equilibrium, an information cascade starts in finite time a.s.

### **Proof**

$$\mathsf{WTS}\,\{q_\infty=\hat{q}_0^+\},\,\mathsf{a.s.}\,\,\exists T<\infty:\forall t\geq T,\,q_t=\hat{q}_0^+.$$

Suppose not. Then wp>0, 
$$\{q_{\infty} = \hat{q}_0^+\} \cap_{t \in \mathbb{N}} \{q_t > \hat{q}_0^+\}.$$

$$\text{On } \{q_{\infty} = \hat{q}^+_0\} \cap_{t \in \mathbb{N}} \{q_t > \hat{q}^+_0\}, \text{ a.s. } \forall \epsilon > 0, \exists T < \infty : \forall t \geq T, \ q_t \in (\hat{q}^+_0, \hat{q}^+_0 + \epsilon).$$

Take  $\varepsilon$  small such that  $\forall q_t \in (\hat{q}_0^+, \hat{q}_0^+ + \varepsilon)$ ,  $a_t = 1 \iff s_t = N_S$ .

$$\begin{array}{l} a_t = \mathbf{1} \implies q_{t+1} = \frac{q_t P_S(N_S|H)}{q_t P_S(N_S|H) + (1-q_t) P_S(N_S|L)} > q_t \text{ and, since } q_t \text{ is martingale,} \\ a_t = \mathbf{0} \implies q_{t+1} = \frac{q_t (1-P_S(N_S|H))}{q_t (1-P_S(N_S|H)) + (1-q_t) (1-P_S(N_S|L))} < q_t. \end{array}$$

Can take 
$$\epsilon > 0$$
 small enough s.t. also 
$$\frac{(\hat{q}_0^{\dagger} + \epsilon)(1 - P_S(N_S|H))}{(\hat{q}_n^{\dagger} + \epsilon)(1 - P_S(N_S|H)) + (1 - (\hat{q}_n^{\dagger} + \epsilon))(1 - P_S(N_S|L))} < \hat{q}_0^{\dagger}.$$

And can take 
$$\varepsilon > 0$$
 small enough s.t. also  $\frac{\hat{q}_0^+ P_S(N_S|H)}{\hat{q}_0^+ P_S(N_S|H) + (1-\hat{q}_0^+) P_S(N_S|L)} > \hat{q}_0^+ + \varepsilon$ .

Hence, for small enough  $\epsilon$ ,  $q_t \in (\hat{q}_0^+, \hat{q}_0^+ + \epsilon) \implies q_{t+1} \notin (\hat{q}_0^+, \hat{q}_0^+ + \epsilon)$ . Contradiction.

#### Done!

# Wrong Cascades Happen

#### Theorem

In any equilibrium, a wrong information cascade starts in finite time a.s.

It is possible everyone unwittingly just takes the suboptimal action!

This is despite the fact every period new information arives.

Markets can get it very wrong due to informational externalities.

# Wrong Cascades Happen

#### **Theorem**

In any equilibrium, for  $p_0 \notin \mathcal{J}$ , a wrong information cascade starts in finite time a.s.

### **Proof**

Consider  $\theta = H$ ; symmetric for  $\theta = L$ .

Since  $q_t \in \mathcal{J}^r \implies q_{t+1} = q_t$  and  $\mathcal{J}^r = [\mathbf{0}, \hat{q}_0^+] \cup (\hat{q}_1^+, \mathbf{1}]$ , then  $q_t$  is effectively bounded and so is  $l(q_t)$ .

Let 
$$w^H := \mathbb{P}(q_\infty \in \mathcal{J}_0^r \mid \theta = H)$$
.

$$I(q_{\infty}) < \infty \text{ a.s. } \Longrightarrow \mathbb{E}[I(q_{\infty}) \mid \theta = H, q_{\infty} \in J_a^r] < \infty.$$

Let  $\tau := \inf\{t \geq \mathbf{0} \mid q_t \in \mathcal{J}'\}$ . Note  $q_{\infty} = q_{\tau}$ . By Doob's optional stopping theorem,  $I(p_0) = \mathbb{E}[I(q_{\tau})|\theta = H] = w^H \mathbb{E}[I(q_{\tau})|\theta = H, q_{\tau} \in \mathcal{J}'_0] + (1 - w^H)\mathbb{E}[I(q_{\tau})|\theta = H, q_{\tau} \in \mathcal{J}'_1]$ .

Hence: 
$$w^H = \frac{I(p_0) - \mathbb{E}[I(q_\tau)|\theta = H, q_\tau \in \mathcal{J}_1]}{\mathbb{E}[I(q_\tau)|\theta = H, q_\tau \in \mathcal{J}_0] - I(p_0)} > 0.$$

## Robustness of Cascades

Smith and Sørensen (2000 Ecta) ask how robust are the results

'Crazy' action/person; need to 'ignore deviation'.

Agent with different preferences; others may still learn from their actions.

Agent has more precise information; contrarian action.

## Robustness of Cascades

### 'Crazy type' $m \in A$ :

Always plays action m. (e.g., preference s.t. strictly dominant to play m)

Arrives with probability  $K \kappa_m \geq 0$ , where  $\kappa_m, K \in [0, 1]$  and  $\sum_{m \in \mathcal{A}} \kappa_m = 1$ .

## **'Rational type'** $r \in R = \{1, ..., N_R\}$ :

Type r determines preference  $u(a, \theta, r)$ .

Arrives with probability  $(1 - K)\rho_r > 0$ .

 $N_A^r$  non-weakly dominated actions.

Order actions  $a_1^r, ..., a_{N_{\Delta}^r}$  s.t.  $u(a_1^r, H, r) < ... < u(a_{N_{\Delta}^r}, H, r)$ .

 $I_a^r$ ,  $J_a^r$ , etc. all defined the same.

Cascade from t if  $q_t \in J := \bigcap_r J^r$ . Common knowledge we're in a cascade.

Limit cascade  $q_{\infty} \in J$ .

## Robustness of Cascades

#### **Private Interim Beliefs**:

```
Private beliefs bounded if 0 < \underline{b} < \overline{b} < 1; unbounded if 0 = \underline{b} < \overline{b} = 1. (Recall \overline{b} = \sup \sup P_P and \underline{b} = \inf \sup P_P.)
```

 $P_P^{\theta}(\cdot) = P_P(\cdot \mid \theta)$ . Assume  $P_P^H$  and  $P_P^L$  mutually abs. continuous (no fully-revealing signals wp> **0**).

Immediate that with unbounded private beliefs,  $J_{a_1^r}^r = \{0\}$  and  $J_{a_{N_A}^r}^r = \{1\}$ , and for other actions  $J_a^r = \emptyset$ .

## Robust and Less Robust Results

### **Theorem**

A fully wrong information cascade a.s. never starts.

Condition on  $\theta = H$ .

- (a) With a single 'rational type', a not-fully-wrong limit cascade occurs a.s.
- (b) With a single 'rational type' and unbounded private beliefs there is complete learning:  $q_t \to \mathbf{1}_{\{\theta=H\}}$  a.s.
- (c) With bounded private beliefs and  $q_t \notin J$ , a wrong limit cascade happens with positive probability.
- (d) Letting the support of private beliefs  $[\underline{b}_n, \overline{b}_n] \to [0, 1]$ , the chance of a wrong limit cascade vanishes (continuity at the limit).

### Robust and Less Robust Results

#### Cascade a.s. in finite time artefact of finite S

For generic atomless  $P_P$ , prob contrarian agent becomes vanishingly small when public belief tends to cascade set.

Hence, from martingale property, variation in public belief converges to zero Public belief never actually reaches the cascade set. (see Chamley 2010 Ch. 4.2.)

#### **Herds vs Cascades**

Limit cascade implies herd (with finite A).

Can have herd without a cascade being triggered.

Herd is triggered a.s. in finite time.

### **Fragile Herds**

Cascades tend to be fragile wrt small shocks.

## Overview

- Social Learning: Checking Neighbours' Crop Yields
- Warm-Up: Gaussian-Quadratic Model
- Information Cascades
- 4. Further Topics

## **Further Topics**

Social Non-Learning, See Kartik, Lee, Liu, and Rappoport (2024 Ecta).

Binary state: Except when private beliefs are unbounded, learning is not complete and may herd on wrong action.

Multiple states: unbounded beliefs okay; weaker condition on info + condition on preferences is sufficient

(intermediate preferences and subexponential location-shift information).

### **Heterogeneous Precision and Order of Moves**

Ottaviani and Sørensen (2001 JPubE): Social learning + reputation concerns. Study order of moves.

Holding aggregate precision fixed, better to have it concentrated in single person.

The anti-seniority rule not necessarily optimal.

Increasing the precision of agent can harm.

## **Limited Observability**

Random sample of past actions (Smith and Sørensen, 2020). Smaller samples can improve efficiency (Chamley 2020, Ch. 5).

Ability to observe others' actions/payoffs may cause insufficient exploration. E.g., favourable reviews deter exploration (Acemoglu, Makhdoumi, Malekian, and Ozdaglar (2022 Ecta)).

## **Further Topics**

**Endogeneous Timing** in BHW model often just expands set equilibria.

See Chamley and Gale (1994 Ecta), Chamley (2010 Ch. 6).

See also Gul and Lundholm (1995 JPE), Caplin and Leahy (1994 AER).

**Overconfidence** can help (Bernardo and Welch (2001 JEMS); Arieli, Babichenko, Müller, Pourbabaee, and Tamuz (2024)).

Correlation neglect can harm (Eyster and Rabin (2010 AER)).

### Non-EU Maximising

Uncertainty wrt others' signal precision and uncertainty aversion can deter breaking cascades even in unbounded private beliefs case (Chen (2025 AER)).

28

## **Further Topics**

### **Costly Info**

Ali (2018 JET): Flexible info.

Complete learning with finite A and costly signals iff  $\mathbb{P}(\cos t \leq \epsilon) > 0$  for any  $\epsilon > 0$ .

Similar insight to unbounded beliefs. Also similar result if agents pay search costs for history (Mueller-Frank Pai (2016 AEJMicro)).

### **Changing State**

Levy, Peski, and Vieille (2024 Ecta).

Population each period. Agents sample actions from previous period and can acquire costly private signal.

At eqm, always acquire private signal and learning incomplete even with very precise signals.

(See also Dasaratha, Golub, and Hak (2023 REStud) for changing state + networks)

More: Social learning in markets, networks, contracting, ...

See Bikhchandani, Hirshleifer, Tamuz, and Welch (2024 JEL).

# Social Learning

Duarte Gonçalves
University College London

Topics in Economic Theory